


Houndstongue

Colorado Dept. of
Agriculture,
Conservation
Services Division
700 Kipling Street
Suite 4000
Lakewood, CO 80215
303-239-4100



Key ID Points

1. Panicles of reddish-purple flowers with 5 petals and 5 soft, hairy sepals.
2. Velcro-like seeds with 4 nutlets.

Houndstongue Identification and Management



Identification and Impacts

Houndstongue (*Cynoglossum officianale*) is a short lived perennial or biennial forb. It produces rosettes in the first year, and bolts a stout, erect stem that is 1 to 4 feet tall, by mid-summer of the second year. Then it flowers and produces fruit. Flowers are reddish-purple (occasionally white) and droop slightly from densely clustered panicles. The five rounded petals are cupped by five sepals covered with long, soft white hairs. Flowering occurs May to July. The simple leaves are lance or oblong shaped, with a smooth edge and no teeth or lobes. Leaves are alternate, 1 to 12 inches long and 1 to 3 inches wide. The leaf tip is sharply pointed, like a hound's tongue, yet are covered with long-soft white hairs. Leaves often appear dusty and insect-ridden. A thick, dark, woody taproot can reach 3 to 4 feet deep.

Reproduction is solely by seeds. Seeds are 4 prickly teardrop-shaped nutlets, which are packed in a pyramid-shaped receptacle. Most seeds fall close to the parent plant, but the seeds can travel great distances. The seeds have barbs like Velcro, with a hooked tip that clings to animals, clothing and machinery. A mature plant can produce 2,000 seeds. Seed viability is 1 to 3 years. Houndstongue is poisonous. Toxic pyrrolizidine alkaloids in Houndstongue stop liver cells from reproducing. Livestock and

wildlife may live up to six months after ingesting a lethal dose. Though the plant has a distinctive odor that repels animals, it is more palatable when dried. Animals rarely eat it unless it is dried and mixed with hay. Houndstongue's toxicity effects horses and cattle more severely, sheep seem more resistant. Burs will reduce the value of sheep wool if present.

Habitats for Houndstongue are open to shady, moist, disturbed areas, along trails, roadsides, fields, pasture, rangeland, along the edge of forests, sand dunes and ditch banks. Houndstongue prefers moist areas, but often grows on sandy or gravelly alkaline soil up to 9,000 feet elevation. Areas with more than 10% bare ground are particularly vulnerable to Houndstongue invasions.

The key to effective control of Houndstongue is preventing establishment and to prevent seed production. Planting competing and desirable grasses and forbs can be effective. Helping with reestablishment of disturbed sites. An integrated weed management approach can also be successful. Chemical, mechanical, and biological controls can be effective when dealing with Houndstongue. Details on the back of this sheet can help to create a management plan compatible with your site ecology.

Houndstongue is designated as a "List B" species in the Colorado Noxious Weed Act. It is required to be either eradicated, contained, or suppressed depending on the local infestations. For more information visit www.colorado.gov/ag/csd and click on the Noxious Weed Management Program. Or call the State Weed Coordinator at the Colorado Department of Agriculture, Conservation Services Division, 303-239-4100.

Photos: top left Aspen County, CO; all other Kelly Uhing, Colorado Department of Agriculture.

Cynoglossum officianale

**CULTURAL**

Prevent the establishment of new infestations by minimizing disturbance and seed dispersal, eliminating seed production and maintaining healthy native communities. Contact your local Natural Resources Conservation Service for seed mix recommendations. Maintain healthy pastures and prevent bare spots caused by overgrazing.

**BIOLOGICAL**

A root weevil, *Mogulones cruciger*, has been successful in Canada and introduced in Montana, but has not yet been approved for use in Colorado. For more information, contact the Palisade Insectary of the Colorado Department of Agriculture at 970-464-7916.

**MECHANICAL**

Cut or pull plants, and remove entire root crown when plants are in the rosette stage. Remove dense litter layer (up to 4 inches) to stimulate germination of desired plants. To reduce seed production, mow or cut flowering stems before seed nutlets develop, this can significantly reduce seed production.

Integrated Weed Management:

Prevention is the best option when dealing with Houndstongue. Use only certified weed-free hay. If an infestation does occur, reducing the seed production is key in controlling Houndstongue. Chemical, mechanical and the developing biological controls can also be effective management techniques.

Houndstongue

HERBICIDES

NOTE: The following are recommendations for herbicides that can be applied to specific areas. Rates are approximate and based on equipment with an output of 30 gal./acre. **Always read, understand, and follow the label directions. The herbicide label is the LAW!**

HERBICIDE	RATE	APPLICATION TIMING
Metsulfuron Methyl + Chlorsulfuron (Cimarron X-tra)	2.0 oz. product/acre + 0.25% v/v non-ionic surfactant	Apply in spring rosette to early bud growth stages.
Picloram + 2,4-D (Grazon P+D)	4 pints/acre + 0.25% v/v non-ionic surfactant	Apply in spring rosette stage.